

Approximation of functions from $L^p(\omega)_\beta$ by matrix means of their Fourier series

Radosław Kranz, Włodzimierz Lenski and Bogdan Szal

University of Zielona Góra

Faculty of Mathematics, Computer Science and Econometrics

65-516 Zielona Góra, ul. Szafrana 4a, Poland

R.Kranz@wmie.uz.zgora.pl ,

W.Lenski@wmie.uz.zgora.pl ,

B.Szal @wmie.uz.zgora.pl

Abstract

We formulate some special conditions for the integrable functions and moduli of continuity. We give the results on rate of approximation of such functions by matrix means of their Fourier series, where the entries of the rows of the matrix generate the sequences belonging to the classes *MRBVS* and *MHBVS*. We also present some results on norm approximation for functions from the generalized integral Lipschitz classes.

Key words: Rate of approximation, summability of Fourier series, Lipschitz classes

2000 Mathematics Subject Classification: 42A24

1 Introduction

Let L^p ($1 \leq p < \infty$) be the class of all 2π -periodic real-valued functions integrable in the Lebesgue sense with p -th power over $Q = [-\pi, \pi]$ with the seminorm

$$\|f\| := \|f(\cdot)\|_{L^p} = \left(\int_Q |f(t)|^p dt \right)^{1/p} \quad (1)$$

and consider the trigonometric Fourier series

$$Sf(x) := \frac{a_0(f)}{2} + \sum_{\nu=1}^{\infty} (a_\nu(f) \cos \nu x + b_\nu(f) \sin \nu x)$$

with the partial sums $S_k f$.

Let $A := (a_{n,k})$ be an infinite lower triangular matrix of real numbers such that

$$a_{n,k} \geq 0 \text{ when } k = 0, 1, 2, \dots, n, \quad a_{n,k} = 0 \text{ when } k > n,$$

$$\sum_{k=0}^n a_{n,k} = 1, \text{ where } n = 0, 1, 2, \dots,$$

and let the A -transformation of $(S_k f)$ be given by

$$T_{n,A} f(x) := \sum_{k=0}^n a_{n,k} S_k f(x) \quad (n = 0, 1, 2, \dots).$$

Denote, for $m = 0, 1, 2, \dots, n$,

$$A_{n,m} = \sum_{k=0}^m a_{n,k} \quad \text{and} \quad \bar{A}_{n,m} = \sum_{k=m}^n a_{n,k}.$$

We define two classes of sequences (see [4]).

A sequence $c := (c_k)$ of nonnegative numbers tending to zero is called the Rest Bounded Variation Sequence, or briefly $c \in RBVS$, if it has the property

$$\sum_{k=m}^{\infty} |c_k - c_{k+1}| \leq K(c) c_m$$

for all positive integer m , where $K(c)$ is a constant depending only on c .

A sequence $c := (c_k)$ of nonnegative numbers will be called the Head Bounded Variation Sequence, or briefly $c \in HBVS$, if it has the property

$$\sum_{k=0}^{m-1} |c_k - c_{k+1}| \leq K(c) c_m$$

for all positive integer m , or only for all $m \leq n$ if the sequence c has only finite nonzero terms and the last nonzero term is c_n .

Now, we define two another classes of sequences.

Follows by L. Leindler (see [5]) a sequence $c := (c_k)$ of nonnegative numbers tending to zero is called the Mean Rest Bounded Variation Sequence, or briefly $c \in MRBVS$, if it has the property

$$\sum_{k=m}^{\infty} |c_k - c_{k+1}| \leq K(c) \frac{1}{m+1} \sum_{k \geq m/2}^m c_k \quad (2)$$

for all positive integer m .

Similar classes of sequences were considered in the papers [2], [6] and [9].

Analogously, a sequence $c := (c_k)$ of nonnegative numbers will be called the Mean Head Bounded Variation Sequence, or briefly $c \in MHBVS$, if it has the property

$$\sum_{k=0}^{n-m-1} |c_k - c_{k+1}| \leq K(c) \frac{1}{m+1} \sum_{k=n-m}^n c_k, \quad (3)$$

for all positive integer $m < n$, where the sequence c has only finite nonzero terms and the last nonzero term is c_n .

It is clear that (see [8])

$$RBVS \subsetneq MRBVS \quad \text{and} \quad HBVS \subsetneq MHBVS.$$

Consequently, we assume that the sequence $(K(\alpha_n))_{n=0}^\infty$ is bounded, that is, that there exists a constant K such that

$$0 \leq K(\alpha_n) \leq K$$

holds for all n , where $K(\alpha_n)$ denote the sequence of constants appearing in the inequalities (2) or (3) for the sequence $\alpha_n = (a_{n,k})_{k=0}^n$, $n = 0, 1, 2, \dots$

Now we can give the conditions to be used later on. We assume that for all n and $0 \leq m < n$

$$\sum_{k=m}^{n-1} |a_{n,k} - a_{n,k+1}| \leq K \frac{1}{m+1} \sum_{k \geq m/2}^m a_{n,k} \quad (4)$$

and

$$\sum_{k=0}^{n-m-1} |a_{n,k} - a_{n,k+1}| \leq K \frac{1}{m+1} \sum_{k=n-m}^n a_{n,k} \quad (5)$$

hold if $(a_{n,k})_{k=0}^n$ belongs to $MRBVS$ or $MHBVS$, for $n = 1, 2, \dots$, respectively.

As a measure of approximation of f by $T_{n,A}f$ we use the generalized modulus of continuity of f in the space L^p defined for $\beta \geq 0$ by the formula

$$\omega_\beta f(\delta)_{L^p} := \sup_{0 \leq |t| \leq \delta} \left\{ \left| \sin \frac{t}{2} \right|^{\beta p} \int_0^\pi |\varphi_x(t)|^p dx \right\}^{\frac{1}{p}},$$

where

$$\varphi_x(t) := f(x+t) + f(x-t) - 2f(x).$$

It is clear that for $\beta \geq \alpha \geq 0$

$$\omega_\beta f(\delta)_{L^p} \leq \omega_\alpha f(\delta)_{L^p}$$

and it is easily seen that $\omega_0 f(\cdot)_{L^p} = \omega f(\cdot)_{L^p}$ is the classical modulus of smoothness.

In our theorems we shall consider the pointwise deviation $T_{n,A}f(x) - f(x)$. We shall formulate the special conditions for the functions and moduli of continuity. Consequently, we also give some results on norm approximation. We shall use the matrix which the entries of the rows generate the sequences belonging to the classes of sequences $MRBVS$ and $MHBVS$. Thus we essentially extend and improve the earlier results [3, Theorem 2, p. 347] and [7]. Moreover, we prove that from our results the pointwise version of the correct form of the Lal theorem follows.

We shall write $I_1 \ll I_2$ if there exists a positive constant K , sometimes depending on some parameters, such that $I_1 \leq KI_2$.

2 Statement of the results

Let us consider a function ω of modulus of continuity type on the interval $[0, 2\pi]$, i.e. a nondecreasing continuous function having the following properties: $\omega(0) = 0$, $\omega(\delta_1 + \delta_2) \leq \omega(\delta_1) + \omega(\delta_2)$ for any $0 \leq \delta_1 \leq \delta_2 \leq \delta_1 + \delta_2 \leq 2\pi$. It is easy to conclude that the function $\delta^{-1}\omega(\delta)$ is quasi nonincreasing function of δ . Let

$$L^p(\omega)_\beta = \{f \in L^p : \omega_\beta f(\delta)_{L^p} \ll \omega(\delta)\}.$$

It is clear that, for $\beta > \alpha \geq 0$,

$$L^p(\omega)_\alpha \subset L^p(\omega)_\beta.$$

Now, we can formulate our main results on the degrees of pointwise summability. The pointwise nature of our results ensure the assumptions of presented below theorems with $f(x)$ (cf. [1]).

Theorem 1 *Let $f \in L^p$ ($1 < p < \infty$) and let ω satisfy*

$$\sum_{m=1}^n (m+1)^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \left(\frac{|\varphi_x(t)|}{\omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} = O_x \left((n+1)^{\beta+1/p} \right) \quad (6)$$

and

$$\left\{ \int_0^{\frac{2\pi}{n}} \left(\frac{|\varphi_x(t)|}{\omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} = O_x \left((n+1)^{-1/p} \right) \quad (7)$$

with $0 \leq \beta < 1 - \frac{1}{p}$ and $q = \frac{p}{p-1}$. If $(a_{n,k})_{k=0}^n \in MRBVS$, then

$$|T_{n,A}f(x) - f(x)| = O_x \left((n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,k} \omega \left(\frac{\pi}{k+1} \right) \right)$$

and if $(a_{n,k})_{k=0}^n \in MHBVS$, then

$$|T_{n,A}f(x) - f(x)| = O_x \left((n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,n-k} \omega \left(\frac{\pi}{k+1} \right) \right),$$

for considered x .

Supposing a stronger condition than (6) we can obtain a better order of approximation.

Theorem 2 *Let $f \in L^p$ ($1 < p < \infty$) and let ω satisfy*

$$\sum_{m=1}^n (m+1)^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \left(\frac{|\varphi_x(t)|}{\omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} = O_x \left((n+1)^\beta \right) \quad (8)$$

and (7) with $0 \leq \beta < 1 - \frac{1}{p}$ and $q = \frac{p}{p-1}$. If $(a_{n,k})_{k=0}^n \in MRBVS$, then

$$|T_{n,A}f(x) - f(x)| = O_x \left((n+1)^\beta \sum_{k=0}^n a_{n,k} \omega \left(\frac{\pi}{k+1} \right) \right)$$

and if $(a_{n,k})_{k=0}^n \in MHBVS$, then

$$|T_{n,A}f(x) - f(x)| = O_x \left((n+1)^\beta \sum_{k=0}^n a_{n,n-k} \omega \left(\frac{\pi}{k+1} \right) \right),$$

for considered x .

Next, we formulate the result on estimates in the L^p norm of the deviation considered above.

Theorem 3 Let $f \in L^p(\omega)_\beta$ ($1 < p < \infty$) with $0 \leq \beta < 1 - \frac{1}{p}$. If $(a_{n,k})_{k=0}^n \in MRBVS$, then

$$\|T_{n,A}f(\cdot) - f(\cdot)\|_{L^p} = \begin{cases} O \left((n+1)^\beta \sum_{k=0}^n a_{n,k} \omega \left(\frac{\pi}{k+1} \right) \right) & \text{for } \beta > 0, \\ O \left((n+1)^{\frac{1}{p}} \sum_{k=0}^n a_{n,k} \omega \left(\frac{\pi}{k+1} \right) \right) & \text{for } \beta = 0, \end{cases}$$

and if $(a_{n,k})_{k=0}^n \in MHBVS$, then

$$\|T_{n,A}f(\cdot) - f(\cdot)\|_{L^p} = \begin{cases} O \left((n+1)^\beta \sum_{k=0}^n a_{n,n-k} \omega \left(\frac{\pi}{k+1} \right) \right) & \text{for } \beta > 0, \\ O \left((n+1)^{\frac{1}{p}} \sum_{k=0}^n a_{n,n-k} \omega \left(\frac{\pi}{k+1} \right) \right) & \text{for } \beta = 0. \end{cases}$$

Finally, we give some remarks and corollary.

Remark 1 Analyzing the proofs of our Theorems we can deduce that under the additional assumption $A_{n,k} = O \left(\frac{k}{n+1} \right)$ or $\bar{A}_{n,k} = O \left(\frac{k}{n+1} \right)$ we obtain the expression $\omega \left(\frac{\pi}{n+1} \right)$ instead of $\sum_{k=0}^n a_{n,k} \omega \left(\frac{\pi}{k+1} \right)$ or $\sum_{k=0}^n a_{n,n-k} \omega \left(\frac{\pi}{k+1} \right)$, respectively, in the estimates of our deviation $T_{n,A}f - f$.

Remark 2 We can observe that taking $a_{n,k} = \frac{1}{n+1} \sum_{\nu=k}^n \frac{p_{\nu-k}}{P_\nu}$ we obtain the mean considered in [3] and if (p_ν) is monotonic with respect to ν then $(a_{n,k})_{k=0}^n \in MRBVS$. Moreover, under the assumptions of Theorem 1 on a function f , if (p_ν) is nonincreasing sequence such that the condition

$$P_\tau \sum_{\nu=\tau}^n P_\nu^{-1} = O(\tau)$$

holds, then from Remark 1, using Lemma 3, we can obtain the correct form of the result of S. Lal [3].

Corollary 1 *If $f \in Lip(\alpha, p) := \{g \in L^p : \omega g(\delta)_{L^p} \ll \delta^\alpha\}$ ($0 < \alpha \leq 1$, $1 \leq p < \infty$), then $f \in L^p(\omega)_\beta$ with $\omega(\delta) = \delta^{\alpha+\beta}$, where $\alpha + \beta \leq 1$. Taking the Fejer mean, i.e. $a_{n,k} = \frac{1}{n+1}$ for $k = 0, 1, \dots, n$ and $a_{n,k} = 0$ for $k > n$, we get from Theorem 3, by Remark 1 with $\alpha \in (0, 1)$, the following estimate*

$$\left\| T_{n, (\frac{1}{n+1})} f(\cdot) - f(\cdot) \right\|_{L^p} = O\left(\frac{1}{(n+1)^\alpha}\right)$$

for $\beta \in (0, 1 - \frac{1}{p})$, where $1 < p < \infty$. So, we obtain the best order of approximation on wider class of functions.

We can also deduce further remarks.

Remark 3 *Analyzing the proof of Theorem 1 we can deduce that taking the assumption $(a_{n,k})_{k=0}^n \in RBVS$ or $(a_{n,k})_{k=0}^n \in HBVS$ instead of $(a_{n,k})_{k=0}^n \in MRBVS$ or $(a_{n,k})_{k=0}^n \in MHBVS$, respectively, we obtain the results from the paper [7].*

Remark 4 *In the case $p \geq 1$ (specially if $p = 1$) we can suppose that the expression $t^{-\beta} \omega(t)$ is nondecreasing in t instead of the assumption $\beta < 1 - \frac{1}{p}$.*

3 Auxiliary results

We begin this section by some notations following A. Zygmund ([10]). It is clear that

$$S_k f(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x+t) D_k(t) dt$$

and

$$T_{n,A} f(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x+t) \sum_{k=0}^n a_{n,k} D_k(t) dt,$$

where

$$D_k(t) = \frac{1}{2} + \sum_{\nu=1}^k \cos \nu t = \begin{cases} \frac{\sin \frac{(2k+1)t}{2}}{2 \sin \frac{t}{2}} & \text{for } t \neq 2l\pi, \\ k + \frac{1}{2} & \text{for } t = 2l\pi. \end{cases}$$

Hence

$$T_{n,A} f(x) - f(x) = \frac{1}{\pi} \int_0^\pi \varphi_x(t) \sum_{k=0}^n a_{n,k} D_k(t) dt.$$

Next, we present the known estimates for the Dirichlet kernel.

Lemma 1 [10] *If $0 < |t| \leq \pi$ then*

$$|D_k(t)| \leq \frac{\pi}{|t|}$$

and, for any real t , we have

$$|D_k(t)| \leq k + 1.$$

More complicated estimates we give with proofs.

Lemma 2 (cf. [4, 7, 8]) If $(a_{n,k})_{k=0}^n \in MHBVS$, then

$$\left| \sum_{k=0}^n a_{n,k} D_k(t) \right| = O(t^{-1} \overline{A}_{n,n-2\tau}),$$

and if $(a_{n,k})_{k=0}^n \in MRBVS$, then

$$\left| \sum_{k=0}^n a_{n,k} D_k(t) \right| = O(t^{-1} A_{n,\tau}),$$

for $\frac{2\pi}{n} \leq t \leq \pi$ ($n = 2, 3, \dots$), where $\tau = [\pi/t]$.

Proof. Using partial summation

$$\begin{aligned} & \left| \sum_{k=0}^n a_{n,k} \sin \frac{(2k+1)t}{2} \sin \frac{t}{2} \right| \\ & \leq tA_{n,\tau} + \frac{1}{2} \sum_{k=\tau-1}^{n-1} |a_{n,k} - a_{n,k+1}| \left| \sum_{l=\tau-1}^k 2 \sin \frac{(2l+1)t}{2} \sin \frac{t}{2} \right| \\ & \quad + \frac{1}{2} a_{n,n} \left| \sum_{l=\tau-1}^n 2 \sin \frac{(2l+1)t}{2} \sin \frac{t}{2} \right| \\ & \leq tA_{n,\tau} + \frac{1}{2} \sum_{k=\tau-1}^{n-1} |a_{n,k} - a_{n,k+1}| \left| \sum_{l=\tau-1}^k (\cos lt - \cos(l+1)t) \right| \\ & \quad + \frac{1}{2} a_{n,n} \left| \sum_{l=\tau-1}^n (\cos lt - \cos(l+1)t) \right| \\ & \leq tA_{n,\tau} + \frac{1}{2} \sum_{k=\tau-1}^{n-1} |a_{n,k} - a_{n,k+1}| |\cos(\tau-1)t - \cos(k+1)t| \\ & \quad + \frac{1}{2} a_{n,n} |\cos(\tau-1)t - \cos(n+1)t| \\ & \leq tA_{n,\tau} + \sum_{k=\tau}^{n-1} |a_{n,k} - a_{n,k+1}| + a_{n,n} \end{aligned}$$

or

$$\begin{aligned}
& \left| \sum_{k=0}^n a_{n,k} \sin \frac{(2k+1)t}{2} \sin \frac{t}{2} \right| \\
& \leq t\bar{A}_{n,n-\tau} + \frac{1}{2} \sum_{k=0}^{n-\tau-1} |a_{n,k} - a_{n,k+1}| \left| \sum_{l=0}^k 2 \sin \frac{(2l+1)t}{2} \sin \frac{t}{2} \right| \\
& \quad + \frac{1}{2} a_{n,n-\tau} \left| \sum_{l=0}^{n-\tau} 2 \sin \frac{(2l+1)t}{2} \sin \frac{t}{2} \right| + a_{n,n-\tau} \\
& \leq t\bar{A}_{n,n-\tau} + \sum_{k=0}^{n-\tau-1} |a_{n,k} - a_{n,k+1}| + 2a_{n,n-\tau}.
\end{aligned}$$

Because $(a_{n,k})_{k=0}^n \in MRBVS$ we have

$$\begin{aligned}
a_{n,s+1} - a_{n,m} & \leq |a_{n,m} - a_{n,s+1}| \leq \sum_{k=m}^s |a_{n,k} - a_{n,k+1}| \\
& \leq \sum_{k=r}^{n-1} |a_{n,k} - a_{n,k+1}| \ll \frac{1}{r+1} \sum_{k \geq r/2}^r a_{n,k} \quad (0 \leq r \leq m \leq s < n),
\end{aligned}$$

whence

$$a_{n,s+1} \ll a_{n,m} + \frac{1}{r+1} \sum_{k \geq r/2}^r a_{n,k} \quad (0 \leq r \leq m \leq s < n)$$

and therefore

$$\begin{aligned}
& \left| \sum_{k=0}^n a_{n,k} \sin \frac{(2k+1)t}{2} \sin \frac{t}{2} \right| \\
& \ll tA_{n,\tau} + \frac{1}{\tau+1} \sum_{k \geq \tau/2}^{\tau} a_{n,k} + \frac{1}{\tau+1} \sum_{k \geq \tau/2}^{\tau} a_{n,n} \\
& \ll tA_{n,\tau} + \frac{1}{\tau+1} \sum_{k \geq \tau/2}^{\tau} a_{n,k} + \frac{1}{\tau+1} \sum_{m \geq \tau/2}^{\tau} \left(a_{n,m} + \frac{1}{\tau+1} \sum_{k \geq \tau/4}^{\tau/2} a_{n,k} \right) \ll tA_{n,\tau}.
\end{aligned}$$

Analogously, the relation $(a_{n,k})_{k=0}^n \in MHBVS$ implies

$$\begin{aligned}
a_{n,s} - a_{n,m} & \leq |a_{n,m} - a_{n,s}| \leq \sum_{k=m}^{s-1} |a_{n,k} - a_{n,k+1}| \\
& \leq \sum_{k=0}^{r-1} |a_{n,k} - a_{n,k+1}| \\
& \ll \frac{1}{n-r+1} \sum_{k=r}^n a_{n,k} \quad (0 \leq m < s \leq r \leq n)
\end{aligned}$$

and

$$a_{n,s} \ll a_{n,m} + \frac{1}{n-r+1} \sum_{k=r}^n a_{n,k} \quad (0 \leq m < s \leq r \leq n),$$

whence

$$\begin{aligned} & \left| \sum_{k=0}^n a_{n,k} \sin \frac{(2k+1)t}{2} \sin \frac{t}{2} \right| \\ & \ll t \bar{A}_{n,n-\tau} + \frac{1}{\tau+1} \sum_{k=n-\tau}^n a_{n,k} + a_{n,n-\tau} \frac{1}{\tau} \sum_{m=n-2\tau}^{n-\tau-1} 1 \\ & \ll t \bar{A}_{n,n-\tau} + \frac{1}{\tau} \sum_{m=n-2\tau}^{n-\tau-1} \left(a_{n,m} + \frac{1}{\tau+1} \sum_{k=n-\tau}^n a_{n,k} \right) \\ & \leq 2t \bar{A}_{n,n-\tau} + \frac{1}{\tau} \sum_{k=n-2\tau}^{n-\tau-1} a_{n,k} \ll t \bar{A}_{n,n-2\tau} \quad \left(\tau \leq \frac{n}{2} \right). \end{aligned}$$

Thus our proof is complete. ■

Lemma 3 Let $f \in L^p$ ($1 < p < \infty$), $\beta \geq 0$ and $0 \leq \gamma < \beta + \frac{1}{p}$. If

$$\left\{ \int_{\frac{\pi}{n+1}}^{\pi} \left(\frac{|\varphi_x(t)|}{t^\gamma \omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{\frac{1}{p}} = O_x((n+1)^\gamma), \quad (9)$$

then

$$\sum_{m=1}^n (m+1)^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \left(\frac{|\varphi_x(t)|}{\omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} = O_x((n+1)^{\beta+1/p}).$$

Proof. Let (9) holds. Then, using the Hölder inequality $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$, we have

$$\begin{aligned} & \sum_{m=1}^n (m+1)^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \left(\frac{|\varphi_x(t)|}{\omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} \\ & \ll \sum_{m=1}^n (m+1)^{\beta-\gamma+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \left(\frac{|\varphi_x(t)|}{t^\gamma \omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} \\ & \leq \left\{ \sum_{m=1}^n \left((m+1)^{\beta-\gamma+1-\frac{2}{q}} \right)^q \right\}^{\frac{1}{q}} \left\{ \sum_{m=1}^n \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \left(\frac{|\varphi_x(t)|}{t^\gamma \omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{1/p} \\ & \leq \left\{ \sum_{m=1}^n (m+1)^{\beta q - \gamma q + q - 2} \right\}^{\frac{1}{q}} \left\{ \int_{\frac{\pi}{n+1}}^{\pi} \left(\frac{|\varphi_x(t)|}{t^\gamma \omega(t)} \right)^p \sin^{\beta p} \frac{t}{2} dt \right\}^{\frac{1}{p}} \\ & \ll (n+1)^{\beta+1/p} \end{aligned}$$

when $0 \leq \gamma < \beta + \frac{1}{p}$. ■

4 Proofs of the results

4.1 Proof of Theorem 1

Let

$$\begin{aligned} T_{n,A}f(x) - f(x) &= \frac{1}{\pi} \int_0^{\frac{2\pi}{n}} \varphi_x(t) \sum_{k=0}^n a_{n,k} D_k(t) dt \\ &\quad + \frac{1}{\pi} \int_{\frac{2\pi}{n}}^{\pi} \varphi_x(t) \sum_{k=0}^n a_{n,k} D_k(t) dt \\ &= I_1 + I_2, \end{aligned}$$

where $n \geq 2$. Then

$$|T_{n,A}f(x) - f(x)| \leq |I_1| + |I_2|.$$

By the Hölder inequality $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$, Lemma 1 and (7), for $\beta < 1 - \frac{1}{p}$,

$$\begin{aligned} |I_1| &\leq \frac{(n+1)}{\pi} \int_0^{\frac{2\pi}{n}} |\varphi_x(t)| dt \\ &\leq \frac{(n+1)}{\pi} \left\{ \int_0^{\frac{2\pi}{n}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_0^{\frac{2\pi}{n}} \left[\frac{\omega(t)}{\sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\ &\ll (n+1)^{1-\frac{1}{p}} \omega\left(\frac{2\pi}{n}\right) \left\{ \int_0^{\frac{2\pi}{n}} \left[\frac{1}{t^\beta} \right]^q dt \right\}^{\frac{1}{q}} \ll (n+1)^\beta \omega\left(\frac{\pi}{n+1}\right). \end{aligned}$$

Suppose that $(a_{n,k})_{k=0}^n \in MRBVS$. By the Hölder inequality $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$, Lemma 2 and (6) for an even n

$$\begin{aligned} |I_2| &\ll \int_{\frac{2\pi}{n}}^{\pi} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt = \sum_{m=1}^{\frac{n}{2}-1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt \\ &\leq \sum_{m=1}^{\frac{n}{2}-1} \sum_{k=0}^{m+1} a_{n,k} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} dt \leq \sum_{m=2}^{\frac{n}{2}} \sum_{k=0}^m a_{n,k} \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \frac{|\varphi_x(t)|}{t} dt \\ &= \sum_{k=2}^{\frac{n}{2}} a_{n,k} \sum_{m=k}^{\frac{n}{2}} \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \frac{|\varphi_x(t)|}{t} dt + \sum_{k=0}^1 a_{n,k} \sum_{m=2}^{\frac{n}{2}} \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \frac{|\varphi_x(t)|}{t} dt \end{aligned}$$

$$\begin{aligned}
&\leq \sum_{k=2}^{\frac{n}{2}} a_{n,k} \sum_{m=k}^{\frac{n}{2}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\
&\quad + \sum_{k=0}^1 a_{n,k} \sum_{m=2}^{\frac{n}{2}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\
&\leq \sum_{k=2}^{\frac{n}{2}} a_{n,k} \sum_{m=k}^{\frac{n}{2}} \omega\left(\frac{\pi}{m-1}\right) \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} t^{(-1-\beta)q} dt \right\}^{\frac{1}{q}} \\
&\quad + \sum_{k=0}^1 a_{n,k} \sum_{m=2}^{\frac{n}{2}} \omega\left(\frac{\pi}{m-1}\right) \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} t^{(-1-\beta)q} dt \right\}^{\frac{1}{q}} \\
&\ll \sum_{k=0}^{\frac{n}{2}} a_{n,k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\frac{n}{2}} m^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \\
&\ll (n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,k} \omega\left(\frac{\pi}{k+1}\right),
\end{aligned}$$

and for an odd n ($n > 2$)

$$\begin{aligned}
|I_2| &\ll \int_{\frac{2\pi}{n}}^{\pi} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt \\
&= \sum_{m=1}^{\left[\frac{n}{2}\right]-1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt + \int_{\frac{2\pi}{n}}^{\left[\frac{\pi}{2}\right]} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt = \sum_1 + \sum_2,
\end{aligned}$$

the first sum we can estimate similarly like for an even n and therefore

$$\begin{aligned}
\sum_1 &\ll \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\left[\frac{n}{2}\right]} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} t^{(-1-\beta)q} dt \right\}^{\frac{1}{q}} \\
&\ll (n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,k} \omega\left(\frac{\pi}{k+1}\right).
\end{aligned}$$

For the second sum we have

$$\sum_2 \leq \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,k} \int_{\frac{2\pi}{n}}^{\left[\frac{\pi}{2}\right]} \frac{|\varphi_x(t)|}{t} dt \leq \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,k} \int_{\left[\frac{n}{2}\right]+1}^{\left[\frac{\pi}{2}\right]} \frac{|\varphi_x(t)|}{t} dt$$

$$\begin{aligned}
&\leq \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\
&\ll \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \omega \left(\frac{\pi}{n+1} \right) \left(\lfloor \frac{n}{2} \rfloor + 1 \right)^{1+\beta-\frac{2}{q}} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \\
&\ll \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \left(\lfloor \frac{n}{2} \rfloor + 1 \right)^{\beta+\frac{1}{p}} \omega \left(\frac{\pi}{n+1} \right) \ll (n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,k} \omega \left(\frac{\pi}{k+1} \right).
\end{aligned}$$

Consequently, let $(a_{n,k})_{k=0}^n \in MHBVS$. By the Hölder inequality $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$, Lemma 2 and (6) for an even n

$$\begin{aligned}
|I_2| &\ll \int_{\frac{2\pi}{n}}^{\pi} \frac{|\varphi_x(t)|}{t} \sum_{k=n-2\tau}^n a_{n,k} dt = \sum_{m=1}^{\frac{n}{2}-1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,n-2k} dt \\
&\leq \sum_{k=2}^{\frac{n}{2}} a_{n,n-2k} \sum_{m=k}^{\frac{n}{2}} \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \frac{|\varphi_x(t)|}{t} dt + \sum_{k=0}^1 a_{n,n-2k} \sum_{m=2}^{\frac{n}{2}} \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \frac{|\varphi_x(t)|}{t} dt \\
&\leq \sum_{k=2}^{\frac{n}{2}} a_{n,n-2k} \sum_{m=k}^{\frac{n}{2}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\
&\quad + \sum_{k=0}^1 a_{n,n-2k} \sum_{m=2}^{\frac{n}{2}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\
&\ll \sum_{k=0}^{\frac{n}{2}} a_{n,n-2k} \omega \left(\frac{\pi}{k+1} \right) \sum_{m=2}^{\frac{n}{2}} m^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \\
&\ll n^{\beta+\frac{1}{p}} \sum_{k=0}^{\frac{n}{2}} a_{n,n-2k} \omega \left(\frac{\pi}{k+1} \right) \leq (n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,n-k} \omega \left(\frac{\pi}{k+1} \right),
\end{aligned}$$

and for an odd n ($n > 2$)

$$\begin{aligned}
|I_2| &\ll \int_{\frac{2\pi}{n}}^{\pi} \frac{|\varphi_x(t)|}{t} \sum_{k=n-2\tau}^n a_{n,k} dt \\
&= \sum_{m=1}^{\lfloor \frac{n}{2} \rfloor - 1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,n-2k} dt + \int_{\frac{2\pi}{n}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,n-2k} dt = \sum_1 + \sum_2.
\end{aligned}$$

The first sum we can estimate similarly like above

$$\begin{aligned} \sum_1 &\ll \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,n-2k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\lfloor \frac{n}{2} \rfloor} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} t^{(-1-\beta)q} dt \right\}^{\frac{1}{q}} \\ &\ll (n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,n-k} \omega\left(\frac{\pi}{k+1}\right). \end{aligned}$$

For the second sum we have

$$\begin{aligned} \sum_2 &\leq \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,n-2k} \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \frac{|\varphi_x(t)|}{t} dt \\ &\leq \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,n-2k} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\ &\ll \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,n-2k} \left(\left\lfloor \frac{n}{2} \right\rfloor + 1 \right)^{\beta+\frac{1}{p}} \omega\left(\frac{\pi}{n+1}\right) \ll (n+1)^{\beta+\frac{1}{p}} \sum_{k=0}^n a_{n,n-k} \omega\left(\frac{\pi}{k+1}\right). \end{aligned}$$

Collecting these estimates we obtain the desired result. ■

4.2 Proof of Theorem 2

Similarly to the proof of Theorem 1, we have

$$|T_{n,A}f(x) - f(x)| \leq |I_1| + |I_2|.$$

and for $\beta < 1 - \frac{1}{p}$,

$$|I_1| \ll (n+1)^\beta \omega\left(\frac{\pi}{n+1}\right).$$

Suppose that $(a_{n,k})_{k=0}^n \in MRBVS$. By the Hölder inequality $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$, Lemma 2 and (8) for an even n

$$\begin{aligned} |I_2| &\ll \int_{\frac{2\pi}{n}}^{\pi} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt = \sum_{m=1}^{\frac{n}{2}-1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt \\ &\ll \sum_{k=0}^{\frac{n}{2}} a_{n,k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\frac{n}{2}} m^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \\ &\ll (n+1)^\beta \sum_{k=0}^n a_{n,k} \omega\left(\frac{\pi}{k+1}\right), \end{aligned}$$

and for an odd n ($n > 2$)

$$|I_2| \ll \sum_{m=1}^{\lfloor \frac{n}{2} \rfloor - 1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt + \int_{\frac{2\pi}{n}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,k} dt = \sum_1 + \sum_2.$$

For the first sum we have

$$\begin{aligned} \sum_1 &\ll \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\lfloor \frac{n}{2} \rfloor} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} t^{(-1-\beta)q} dt \right\}^{\frac{1}{q}} \\ &\ll (n+1)^\beta \sum_{k=0}^n a_{n,k} \omega\left(\frac{\pi}{k+1}\right) \end{aligned}$$

and for the second one

$$\begin{aligned} \sum_2 &\leq \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \frac{|\varphi_x(t)|}{t} dt \\ &\leq \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\ &\ll \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,k} \omega\left(\frac{\pi}{n+1}\right) \left(\lfloor \frac{n}{2} \rfloor + 1\right)^{1+\beta-\frac{2}{q}} \left\{ \int_{\frac{\pi}{\lfloor \frac{n}{2} \rfloor + 1}}^{\frac{\pi}{\lfloor \frac{n}{2} \rfloor}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \\ &\ll (n+1)^\beta \sum_{k=0}^n a_{n,k} \omega\left(\frac{\pi}{k+1}\right). \end{aligned}$$

Consequently, let $(a_{n,k})_{k=0}^n \in MHBVS$. By the Hölder inequality $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$, Lemma 2 and (8) for an even n

$$\begin{aligned} |I_2| &\ll \int_{\frac{2\pi}{n}}^{\pi} \frac{|\varphi_x(t)|}{t} \sum_{k=n-2\tau}^n a_{n,k} dt = \sum_{m=1}^{\frac{n}{2}-1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,n-2k} dt \\ &\ll \sum_{k=0}^{\frac{n}{2}} a_{n,n-2k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\frac{n}{2}} m^{\beta+1-\frac{2}{q}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \\ &\ll n^\beta \sum_{k=0}^{\frac{n}{2}} a_{n,n-2k} \omega\left(\frac{\pi}{k+1}\right) \ll (n+1)^\beta \sum_{k=0}^n a_{n,n-k} \omega\left(\frac{\pi}{k+1}\right), \end{aligned}$$

and for an odd n ($n > 2$)

$$|I_2| \ll \sum_{m=1}^{\left[\frac{n}{2}\right]-1} \int_{\frac{\pi}{m+1}}^{\frac{\pi}{m}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,n-2k} dt + \int_{\frac{2\pi}{n}}^{\frac{\pi}{\left[\frac{n}{2}\right]}} \frac{|\varphi_x(t)|}{t} \sum_{k=0}^{\tau} a_{n,n-2k} dt = \sum_1 + \sum_2.$$

Analogously as above

$$\begin{aligned} \sum_1 &\ll \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,n-2k} \omega\left(\frac{\pi}{k+1}\right) \sum_{m=2}^{\left[\frac{n}{2}\right]} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{m}}^{\frac{\pi}{m-1}} t^{(-1-\beta)q} dt \right\}^{\frac{1}{q}} \\ &\ll (n+1)^\beta \sum_{k=0}^n a_{n,n-k} \omega\left(\frac{\pi}{k+1}\right) \end{aligned}$$

and

$$\begin{aligned} \sum_2 &\leq \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,n-2k} \int_{\frac{\pi}{\left[\frac{n}{2}\right]+1}}^{\frac{\pi}{\left[\frac{n}{2}\right]}} \frac{|\varphi_x(t)|}{t} dt \\ &\leq \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,n-2k} \left\{ \int_{\frac{\pi}{\left[\frac{n}{2}\right]+1}}^{\frac{\pi}{\left[\frac{n}{2}\right]}} \left[\frac{|\varphi_x(t)|}{\omega(t)} \sin^\beta \frac{t}{2} \right]^p dt \right\}^{\frac{1}{p}} \left\{ \int_{\frac{\pi}{\left[\frac{n}{2}\right]+1}}^{\frac{\pi}{\left[\frac{n}{2}\right]}} \left[\frac{\omega(t)}{t \sin^\beta \frac{t}{2}} \right]^q dt \right\}^{\frac{1}{q}} \\ &\ll \sum_{k=0}^{\left[\frac{n}{2}\right]} a_{n,n-2k} \left(\left[\frac{n}{2} \right] + 1 \right)^\beta \omega\left(\frac{\pi}{n+1}\right) \ll (n+1)^\beta \sum_{k=0}^n a_{n,n-k} \omega\left(\frac{\pi}{k+1}\right). \end{aligned}$$

Collecting these estimates we obtain the desired result. ■

4.3 Proofs of Theorems 3

The proof is similar to these above. The expressions in the estimates under the L^p norm with respect to x will be like these on the left hand side of our conditions (7) and (8) or (6) when $\beta > 0$ or $\beta = 0$, respectively. Since $f \in L^p(\omega)_\beta$, the such norm quantities will always have the same orders like these on the right hand side of the mentioned conditions. Therefore the proof follows without any additional assumptions. ■

References

- [1] P. L. Butzer, R. J. Nessel, Fourier analysis and approximation, Birkhäuser verlag Besel und Stuttgart, 1971.
- [2] M. Dyachenko and S. Tikhonov, Convergence of trigonometric series with general monotone coefficients, C. R. Acad. Sci. Paris, 345(3)(2007), 123-126

- [3] S. Lal, Approximation of functions belonging to the generalized Lipschitz Class by $C1Np$ summability method of Fourier series, Appl. Math. Comput., 209 (2009), 346-350.
- [4] L. Leindler, On the degree of approximation of continuous functions, Acta Math. Hungar., 104 (1-2) (2004), 105-113.
- [5] L. Leindler, Integrability conditions pertaining to Orlicz space, J. Inequal. Pure and Appl. Math., 8(2) (2007), Art. 38, 6 pp.
- [6] E. Liflyand and S. Tikhonov, A concept of general monotonicity and applications, Math. Nachr., 284(2011), 1083-1098
- [7] W. Lenski, B. Szal, Approximation of functions belonging to the class $L^p(\omega)_\beta$ by linear operators, Acta Comment. Univ. Tartu. Math., Vol. 13 (2009), 11-24.
- [8] B. Szal, A note on the uniform convergence and boundedness a generalized class of sine series, Commentat. Math., 48 1 (2008), 85-94.
- [9] S.Tikhonov, Trigonometric series with general monotone coefficients, J. Math. Anal. Appl. 326(1)(2007), 721-735
- [10] A. Zygmund, Trigonometric series, Cambridge, 2002.